arXiv:hep-lat/0010046v1 26 Oct 2000

Deconfinement Through Chiral Transition In 2 Flavour QCD [*]

S. Digal, E. Laermann, H. Satz

Fakultät für Physik, Universität Bielefeld,
D-33501 Bielefeld, Germany

We propose that in QCD with dynamical quarks, colour deconfinement occurs when an external field induced by the chiral condensate strongly aligns the Polyakov loop. This effect sets in at the chiral symmetry restoration temperature T_χ and thus makes deconfinement and chiral symmetry restoration coincide. The predicted singular behavior of Polyakov loop susceptibilities at T_χ is shown to be supported by finite temperature lattice calculations.

1. INTRODUCTION

Hot QCD matter undergoes two well-defined phase transitions at zero baryon chemical potential depending on the bare quark mass parameter. For ($m_q = \infty$) there is the deconfinement transition associated with the spontaneous breaking of the global center symmetry $Z_{N_c} \in SU(N_c)$ of the pure $SU(N_c)$ gauge theory. N_c is the number of colours of quarks. The order parameter describing this transition is the expectation value of the Polyakov loop $L(T)$. There exists a critical temperature T_d, with $L(T) = 0 \ \ \forall \ \ T \leq T_d$ and $L(T) > 0 \ \forall \ T > T_d$. This transition is similar to the magnetic transition in Z_N spin theories with the magnetization $m(T)$ being analogous to $L(T)$. The deconfinement transition and the transition in a spin system have same order when the number of colours N_c and the number of components of spin N are equal [1]. The nature of the deconfinement transition changes with N_c, for $N_c = 2$ the transition is second order and for $N_c \geq 3$ it is first order.

On the other hand for $m_q = 0$, there is the chiral transition at the critical temperature T_x associated with the restoration of chiral symmetry which is spontaneously broken at low temperatures. The order parameter describing this transition is the chiral condensate $K \equiv \langle\bar{\psi}\psi\rangle$. The nature of the chiral transition depends on the number of quark flavours N_f. For $N_f = 2$ the transition is second order and first order for $N_f \geq 3$. In the case of a second order chiral transition the critical behaviour of the chiral condensate are described the critical exponents defined in the following.

$$K(T, m_q = 0) \sim (T_x - T)^\beta, \quad T \lesssim T_x \tag{1}$$

while at $T = T_x$, it vanishes as

$$K(T_x = 0, m_q) \sim m_q^{1/\delta} \tag{2}$$

when $m_q \to 0$. The chiral quark mass susceptibility

$$\begin{aligned} x_m^K(T, m_q) = \left(\frac{\partial K}{\partial m_q}\right)_T &\sim \quad \langle(\psi\bar{\psi})^2\rangle - \langle\psi\bar{\psi}\rangle^2 \\ &\sim \quad |T - T_x|^{-\gamma} \end{aligned} \tag{3}$$

diverges at $T = T_x$ for $m_q \to 0$. β, δ, γ, ... are the critical exponents. It is believed that for $N_f = 2$ the chiral transition is in the universality class of $O(4)$ according to symmetry arguments [2], however it is yet to be confirmed by numerical experiments [3].

Both the symmetries of the Lagrangian, the global Z_{N_c} symmetry and the chiral symmetry, are explicitly broken for finite non-zero m_q. The role of m_q for chiral transition is similar to that of an external magnetic field in spin system, so that for large m_q, the temperature variation of the chiral condensate becomes completely smooth, like the magnetization $m(T)$ in spin systems.

It is not clear what role m_q plays for the deconfinement transition. One might consider the inverse bare quark mass to play such a symmetry breaking role for deconfinement, which would imply that for $m_q \to 0$, the temperature variation of the Polyakov loop expectation value would then become smooth [4]. In lattice studies it is found, however, that this is not the case [5]: even for $m_q \to 0$, the Polyakov loop varies sharply with temperature and the corresponding susceptibility $(\langle L^2\rangle - \langle L\rangle^2)$ peaks sharply at the chiral restoration temperature T_x, which is considerably lower than T_d. In somewhat loose terminology, one describes this situation by saying that in QCD for $m_q \to 0$, deconfinement and chiral symmetry restoration coincide.

---

[*] we are grateful for the financial support by the German Science Ministry BMBF under contract No. 06BI902.

To understand the underlying reason for this coincidence of the critical temperature of chiral and deconfinement transitions we propose [6]that the explicit breaking of the Z_N symmetry is due to the effective external field which is inverse of the constituent quark mass m_Q [7]. It is believed that the constituent quark mass depends on the chiral condensate K, however, the exact dependence is not yet known. If the non-Goldstone hadron masses are related to the chiral condensate, e.g., if the nucleon mass is given by [8]

$$M_n \sim 3\ m_Q \sim K^{1/3}, \tag{4}$$

then chiral symmetry restoration with $K \to 0$ gives rise to a sudden large increase of the external field $H \sim 1/m_Q$, forcing the Polyakov loop to take a large non-zero value. We want to argue that it is this effect which causes a sharp variation of the Polyakov loop $L(T)$ at $T_x < T_d$, with the temperature dependence of $L(T)$ governed by that of the chiral condensate $K(T)$, which is in general different from that obtained in pure gauge theory at T_d.

Although we believe that this interplay of chiral symmetry will in general do not depend on the value of N_f and N_c, for simplicity and to compare with lattice results, we consider $N_f = 2$ and $N_c = 3$ [6]. As we have already mentioned in this case the chiral transition is second order in the chiral limit, and deconfinement is first order in the limit of infinite heavy quarks.

As indicated above, we now take the Z_N structure of QCD to be that of Z_N spin theory in an external field H, determined by the constituent quark mass. From Eq. (4), H and hence also the Polyakov loop thus become functions of K: the quark mass dependence of L enters through $K(T, m_q)$, with $L(T, K)$. In spin systems with external field, the magnetization $m(T, H)$ is for $H \neq 0$ an analytic function of T and H. We therefore assume that $L(T, K)$ is for $\infty > K > 0$ an analytic function of T and K, with the quark mass dependence of L entering through $K(T, m_q)$.

Once the effective external field for deconfinement depends on the chiral condensate K, the critical behavior of the chiral condensate will be reflected in the behavior of the Polyakov loop at $T = T_x$. For a first order chiral transition with a jump in the chiral condensate across the chiral transition, there will be a discontinuity in the effective external field giving rise to a discontinuity in the Polyakov loop across the chiral transition point [9]-[11]. To go further one can write down for the variation of Polyakov loop L,

$$dL = \left(\frac{\partial L}{\partial T}\right)_K dT + \left(\frac{\partial L}{\partial K}\right)_T dK. \tag{5}$$

From this we can calculate the variation of the Polyakov loop with respect to m_q and temperature T as follows,

$$x_m^L = \left(\frac{\partial L}{\partial m_q}\right)_T = \left(\frac{\partial L}{\partial K}\right)_T \left(\frac{\partial K}{\partial m_q}\right)_T \tag{6}$$

and

$$x_T^L = \left(\frac{\partial L}{\partial T}\right)_{m_q} = \left(\frac{\partial L}{\partial T}\right)_K + \left(\frac{\partial L}{\partial K}\right)_T \left(\frac{\partial K}{\partial T}\right)_{m_q}. \tag{7}$$

Both these response functions of the Polyakov loop diverge at $T = T_x$ in the chiral limit $m_q = 0$, since the chiral susceptibilities $x_m^K = (\partial K/\partial m_q)_T$ and $x_T^K = (\partial K/\partial T)_{m_q}$ diverge in this limit.

In [5], the chiral susceptibilities x_T^K and x_m^K were studied on an $8^3 \times 4$ lattice for quark masses $m_q a = 0.075,\ 0.0375$ and 0.02. Results for x_T^K and x_m^K are shown in Figs. 1 and 2, respectively, as functions of the effective temperature variable $T \equiv \kappa = 6/g^2$, where g denotes the coupling in the QCD Lagrangian. The peak heights of both these response functions increase with decreasing quark mass, which indicates the divergence in the chiral limit $m_q \to 0$. In Figs. 3 and 4, we show the corresponding results for the Polyakov loop susceptibilities. They also peak sharply, the



Figure 1. The chiral temperature susceptibility x_κ^K as function of the temperature variable $\kappa = 6/g^2$.



Figure 2. The chiral quark mass susceptibility x_m^K as function of the temperature variable $\kappa = 6/g^2$.



Figure 3. The Polyakov loop temperature susceptibility x_κ^L as function of the temperature variable $\kappa = 6/g^2$.



Figure 4. The Polyakov loop quark mass susceptibility x_m^L as function of the temperature variable $\kappa = 6/g^2$.

peak positions coincide with those for the chiral susceptibilities, and here as well the peak height increases with decreasing quark mass. The observed behavior therefore provides clear support for the divergence of the temperature and quark mass derivatives of the Polyakov loop at T_x in the chiral limit.

A direct comparison of the functional behavior, ideally of the relevant critical exponents, for the two cases becomes difficult for two reasons. The chiral susceptibilities do not at present lead to the predicted $O(4)$ exponents; this may indicate that the quark masses are still too large for pure critical behavior. The Polyakov loop susceptibilities contain in addition unknown non-singular factors $(\partial L/\partial T)_K$ and $(\partial L/\partial K)_T$, which will modify the non-singular m_q-dependence relative to that of the chiral susceptibilities. Lattice studies for smaller quark masses would therefore be very helpful.

In summary, we have shown that chiral symmetry restoration, with the resulting sudden change in an effective constituent quark mass, leads to a suddenly increasing external field which aligns the Polyakov loops at $T = T_x < T_d$ and thus produces a strong explicit breaking of the Z_N symmetry of the gauge field part of the Lagrangian. This effect makes chiral symmetry restoration 'coincide' with deconfinement. The resulting predictions for diverging Polyakov loop susceptibilities are found to be well supported by finite temperature lattice calculations for full QCD with two flavours of light quarks.

REFERENCES

1. B. Svetitsky and L. G. Yaffe, Nucl. Phys. B 210 [FS6] (1982) 423.
2. R. Pisarski and F. Wilczek, Phys. Rev. D29 (1984) 339.
3. S. Aoki et al. (JLQCD), Phys. Rev. D 57 (1998) 3910; For a survey, see E. Laermann, Nucl. Phys. B (Proc. Suppl.) 63 A-C (1998) 114.
4. P. Hasenfratz, F. Karsch and I. Stamatescu, Phys. Lett. 133 B (1983) 221.
5. F. Karsch and E. Laermann, Phys. Rev. D 50 (1994) 6954.
6. S. Digal, E. Laermann and H. Satz, hep-ph/0007175, Euro. Phys. J. C, in press.
7. R. V. Gavai, A. Gocksch, and M. Ogilvie, Phys. Rev. Lett. 56 (1986) 815.
8. B. L. Ioffe, Nucl. Phys. B 188 (1981) 317.
9. R. Gupta et al., Phys. Rev. Lett. 57 (1986) 2621.
10. F. Karsch et al., Phys. Lett. 188 B (1987) 353.
11. R. V. Gavai et al. (MTc), Phys. Lett. B 241 (1990) 567.